

Singing above the chorus: cooperative Princess cichlid fish (*Neolamprologus pulcher*) has high pitch

Rachel K. Spinks · Moritz Muschick ·
Walter Salzburger · Hugo F. Gante 

Received: 19 January 2016 / Revised: 10 July 2016 / Accepted: 12 July 2016
© Springer International Publishing Switzerland 2016

Abstract Teleost fishes not only communicate with well-known visual cues, but also olfactory and acoustic signals. Communicating with sound has advantages, as acoustic signals propagate fast, omnidirectionally, around obstacles and over long distances. Heterogeneous environments might favour multimodal communication, especially in socially complex species, as the combination of modalities' strengths helps overcome their individual limitations. Fishes of the ecologically and morphologically diverse family

Cichlidae are known to be vocal. Here we investigated sound production in the socially complex Princess cichlid *Neolamprologus pulcher* from Lake Tanganyika in East Africa. We show that wild and captive *N. pulcher* produce only short-duration, broadband high-frequency sounds (mean: 12 kHz), when stimulated by mirror images. The evolutionary reasons for this “low frequency silencing” are still unclear. In laboratory experiments, *N. pulcher* produced distinct two-pulsed calls mostly, but not exclusively, associated with agonistic displays. Princess cichlids produce these high-frequency sounds both in combination with and independent from visual displays, suggesting that sounds are not a by-product of behavioural displays. Further studies on the hearing abilities of *N. pulcher* are needed to clarify whether the high-frequency sounds are used in intra- or inter-specific communication.

Guest editors: S. Koblmüller, R. C. Albertson, M. J. Genner, K. M. Sefc & T. Takahashi / Advances in Cichlid Research II: Behavior, Ecology and Evolutionary Biology

Electronic supplementary material The online version of this article (doi:[10.1007/s10750-016-2921-5](https://doi.org/10.1007/s10750-016-2921-5)) contains supplementary material, which is available to authorized users.

R. K. Spinks · M. Muschick · W. Salzburger ·
H. F. Gante (✉)
Zoological Institute, University of Basel, Vesalgasse 1,
4051 Basel, Switzerland
e-mail: hugo.gante@unibas.ch

Present Address:

R. K. Spinks
ARC Centre of Excellence for Coral Reef Studies, James Cook
University, Townsville, QLD 4811, Australia

Present Address:

M. Muschick
Department of Fish Ecology & Evolution, EAWAG Centre for
Ecology, Evolution and Biogeochemistry, 6047 Kastanien-
baum, Switzerland

Keywords Acoustic signals · Sound production ·
High-frequency sound · Low-frequency silencing ·
Neolamprologus pulcher · Lake Tanganyika

Introduction

In spite of the long-held view of a silent underwater world, we now know that many teleost fishes produce sounds as part of their normal behavioural repertoire (Lobel et al., 2010). It should come as no surprise that fish ubiquitously use sounds to communicate, as water is a superior acoustic medium, where sound travels

almost five times faster than in air (Fine & Parmentier, 2015). Compared with other signal modalities, auditory signals can have some advantages: they propagate fast and in all directions unlike olfactory cues, in which case the receiver must be downstream from the sender (Fine & Parmentier, 2015); or around obstacles and to longer distances than visual signals, which quickly become attenuated with increasing distance, in low light, or deep water conditions (Lythgoe & Partridge, 1991). For instance, the nocturnal New Zealand bigeye fish (*Pempheris adspersa* Griffin, 1927) produces sounds mainly at night to promote shoal cohesion when visual cues have reduced utility (Radford et al., 2015; van Oosterom et al., 2016).

Nevertheless, long-range auditory signals also present some communicative weaknesses. For instance, fish need to deal with high levels of environmental noise in shallow water habitats (Ladich & Schulz-Mirbach, 2013; Lugli, 2015), and there is the potential for eavesdropping by non-intended receivers, conspecifics or predators (Verzijden et al., 2010; Bradbury & Vehrencamp, 2011; Maruska et al., 2012). The alternate or simultaneous use of signals of different modalities combines their strengths and reduces limitations imposed by the environment on a particular type of signal (Stevens, 2013). Multimodal communication is thus expected to evolve under varied and unstable environments (Munoz & Blumstein, 2012), in particular in gregarious, territorial and socially complex species (Freeberg et al., 2012).

Fish commonly produce sounds in agonistic, reproductive and defensive contexts (Lobel et al., 2010), either in isolation or most often in association with other signal modalities (Ladich, 1990, 1997). Such sounds are usually low-frequency purrs and grunts (40–1000 Hz), but higher frequency clicks and creaks (above 1 kHz) have also been reported (Ladich, 1997; Lobel et al., 2010). A group of fish that has received increasing attention regarding sound production are cichlids. In particular, the cichlid assemblages of the East African Great Lakes are prime models for studying diversification and adaptation due to varied life histories, morphologies and behaviours (Salzburger, 2009; Gante & Salzburger, 2012; Salzburger et al., 2014). While diversity in colour patterns and visual adaptations has long been recognised as a driving force in cichlid evolution (Santos & Salzburger, 2012; Wagner et al., 2012), the description of sound production and hearing abilities have only more

recently gained momentum in spite of a long history of research (Amorim, 2006; Ladich & Fay, 2013).

Here we report on the production of sounds by the Princess cichlid, *Neolamprologus pulcher* (Trewavas & Poll, 1952). This cooperatively breeding species lives in rocky shores of southern Lake Tanganyika, East Africa, home to one of the most diverse freshwater fish adaptive radiations (Muschick et al., 2012; Salzburger et al., 2014) and has become a favourite in studies of animal cooperation (Wong & Balshine, 2011; Zöttl et al., 2013). In *N. pulcher*, each extended family is typically formed by a dominant breeding couple and up to a few dozen subordinate helpers that collectively raise young and defend their territory from other such groups in the colony. Considering the heterogeneous nature of rocky habitats (especially when compared to sandy habitats) and the high social complexity of cooperative breeders, *N. pulcher* is expected to show increased levels of communicative complexity. Indeed, it has been shown that Princess cichlids use a combination of visual and olfactory signals or cues in multiple aspects of their lives, such as individual recognition, territoriality and aggression (Balshine-Earn & Lotem, 1998; Frostman & Sherman, 2004; Le Vin et al., 2010; Kohda et al., 2015; Bachmann et al., 2016). It is thus puzzling that *N. pulcher* have reportedly gone completely silent (Pisanski et al., 2015). In this study, we further investigate the possibility of sound production in this species by examining both captive bred and wild-caught fish, over a much wider range of sound frequencies than before.

Methods

Acoustic recordings of wild-caught *N. pulcher*—field experiments

Recordings of wild *N. pulcher* were conducted in July and August 2013. *Neolamprologus pulcher* from different social groups were captured with gill nets on SCUBA in shallow waters around Kalambo Lodge, Isanga Bay, Zambia, in the south-eastern shore of Lake Tanganyika (8°37'22.1"S, 31°12'03.6"E). Around 20 adult fish were placed together in a concrete pond (1 × 1 × 1 m), with lake water (mean ± SD; 21.1 ± 0.3°C) and without shelters, so territoriality was subdued and aggression levels reduced between individuals, and left to acclimatise for 3 days before the recordings commenced. In

addition, because we used fish of unknown social status, this period should have allowed for a better homogenisation of their physiological state, which might take hours to a few days to occur (White et al., 2002; Parikh et al., 2006; Maruska & Fernald, 2013), reducing a possible impact on behavioural (including vocalisation) differences between individuals (Buchner et al., 2004; Aubin-Horth et al., 2007; Desjardins et al., 2008). Every second day, one-third of the water in the pond was changed with lake water and the fish fed zooplankton. Fish were individually recorded in another concrete pond that was the same size, but only filled to 20 cm depth, with no other fish present. An octagonal arena, with mirror panels (25 × 20 cm) on the inside, was used to elicit behaviours and sounds (Fig. 1A). Mirrors have been successfully used to induce typical agonistic behaviours in African cichlids and fish in general (Rowland, 1999; Dijkstra et al., 2012). Contrary to the use of other live, interacting fish as stimuli, using mirrors has the advantage that sound emitters cannot be mistaken, and because only one individual is recorded at any one time, precise calculation of sound parameters is also facilitated. To prevent the fish from seeing multiple mirror images, a perforated box was placed in the centre of the arena (Fig. 1A). A Teledyne Reson TC4013 hydrophone (Denmark), with a receiving sensitivity of -211 dB re: V/ μ Pa and frequency range of 1 Hz to 170 kHz, was suspended inside the perforated box. Sound was intensified at 500 Hz by an Ultra-SoundGate charge amplifier and then stored and digitalised at 48 kHz (with 16 bit resolution) into Waveform Audio File Format (.wav) by the Marantz PMD670 recorder. Movements were recorded from above with a GoPro Hero 3 camera that was synchronised to the sound recordings. This allowed discarding sounds that had been produced by the fish touching the setup or breaking the water surface. The pond was illuminated with indirect natural daylight and two solar-charged LED lamps. Unlike fluorescent bulbs, LEDs produce negligible levels of low-frequency sound (Rumyantsev et al., 2005).

Individuals were introduced to the experimental arena via a dark box with a sliding door. After a 2-min acclimatisation period, the door of the box was opened, and the box removed as soon as the fish had vacated it. If the fish did not exit right away, the box was lifted slightly to encourage departure. Each fish was recorded for 8 min and then weighed, standard length measured, and sexed by examining the genital papilla. A total of ten (6 males and 4 females) *N. pulcher* were used in the field experiment. Recordings




Fig. 1 Setups used to record sounds produced by *N. pulcher*. In the field experiment (A), an octagonal mirror arena was used inside a concrete pond (1 × 1 × 1 m) and water levelled to just below the mirror panels (25 × 20 cm). In the laboratory experiment (B), one glass mirror (1.9-mm thick; 28 × 22 cm) was placed against a wall of the aquarium (40 × 30 × 25 cm). In both cases, the hydrophone was immersed *circa* 4 cm

of wild fish taken at Lake Tanganyika were first manually inspected for sounds and then filtered with a bandpass at 300 Hz to remove low-frequency background noise. The experiments were done in accordance with the Department of Fisheries, Lake Tanganyika Research Unit, Mpulungu, Zambia.

Acoustic recordings of captive-raised *N. pulcher*—laboratory experiments

Given the recent report of “silent” *N. pulcher* (Pisanski et al., 2015), sound recordings were repeated under laboratory conditions, where a camera could be

placed in lateral view to monitor fish behaviours with greater detail than in the field (Fig. 1B). It also allowed controlling for the effect of captive raising on sound production, since previous studies have used only captive bred individuals.

In order to minimise ambient background noise, acoustic recordings took place in a room with thick concrete walls, with an aquarium (40 × 30 × 25 cm) resting on 2-cm-thick acoustic absorption cotton and placed inside a large (48 × 42 × 32 cm) expanded polystyrene foam box. The inside of the expanded polystyrene foam container, except for the floor, was also covered with acoustic insulation that allowed external sounds to be reflected and internal sounds to be absorbed to reduce reverberation. Four battery-operated LED lamps were placed above the aquarium to provide adequate illumination. The aquarium contained a half terracotta flowerpot to provide shelter for the fish.

First or second generation laboratory-raised *N. pulcher* were used, originating from fish collected at Kalambo Lodge, Isanga Bay, Zambia in Lake Tanganyika. Fish were originally kept in pairs in aquaria with sandy substrate, halved terracotta flowerpots and a motorised sponge filter, and were fed once daily prior to the experiment. Ten sexually mature *N. pulcher* (5 males and 5 females) were then selected and individually recorded in April 2015. A 1.9-mm-thick glass mirror (28 × 22 cm), placed flat against a lateral wall inside the aquarium, was used to induce sound production (Fig. 1B). Fish were gently hand netted from their home aquaria and given 1 h to acclimatise in the experimental setup; however, the mirror was introduced to the aquarium only 2 min before the recording began to prevent the fish becoming accustomed to it. All nearby electrical equipment, including the room lights, were shut off shortly before synchronous video and audio recordings commenced.

We used the same hydrophone, amplifier, recorder and settings as described in the field experiment. In contrast to the field experiment, in the laboratory recordings, we utilised the Raven Pro 1.5 sound analysis software's adaptive broadband filter, with the default settings of a filter order of ten and a least mean squares step size of 0.01, to reduce the likelihood of filtering out potential fish sounds (Bioacoustics Research Program, 2014). Adaptive broadband filtering is useful when the preferred broadband signal is amidst narrowband background noise that could not otherwise be eliminated (Bioacoustics Research Program, 2014). This filter works just like when people

talk in a noisy environment, the continuous surrounding background sounds are recognised but the focus and concentration is on the person's speech, or in this case on the sounds produced by the fish. To diminish distortion of the fish's acoustic signals in the aquarium, the hydrophone was placed within the attenuation distance of where the fish were expected to produce sound (Akamatsu et al., 2002). Behaviour was simultaneously recorded with a Nikon 1 camera with an 11–27.5 mm lens. Each recording session lasted 20 min. Subsequently, fish were weighed, standard length measured, sexed by examination of the external genital papilla and then returned to their home aquarium. Experiments were authorised by the Cantonal Veterinary Office, Basel, Switzerland (permit numbers 2317 & 2356).

Characterisation of *N. pulcher* sounds

Only sounds that showed a clear structure and high signal-to-noise ratio were considered. All sounds were confirmed with the synchronised video footage and if, for example, the fish touched the mirror or turned around quickly, resulting in an incidental sound, or an unexpected background noise occurred, then no measurements were taken at this time. For this reason, we focused on characterising sounds produced by fish only in the laboratory experiment, where behaviours could be unequivocally monitored. Based on the typical social behaviours of *N. pulcher* (Table 1) we noted if a behavioural display was associated with sound. To quantify the acoustic properties of sounds produced by *N. pulcher* in the laboratory, we measured pulse duration, pulse peak frequency, interpulse interval, call duration and pulse rate (Fig. 2). In the field dataset, we focused on pulse duration and pulse peak frequency. In our study, the duration of each pulse is defined as the time in milliseconds (ms) from the onset of a pulse to its end as classified by amplitude of the signal. Pulse peak frequency is the frequency with the maximum power in the pulse. The duration between each pulse, the interpulse interval, is calculated in milliseconds and is the period with only white noise levels of sound between the pulses. The duration of a call, in milliseconds, is measured from the onset of the first pulse to the end of the last pulse and may contain one pulse or many. Call duration is often subjectively measured in fish acoustics literature. We aimed to provide a non-biased, replicable classification by measuring every interpulse interval in the

Table 1 *Neolamprologus pulcher* ethogram illustrates typical social behaviours of the species (adapted from (Sopinka et al., 2009; Pisanski et al., 2015))

Behaviour	Description
Non-aggressive & social	
Quiver	Fish quivers to mirror; the whole body trembles
Soft touch	Fish nips or softly makes contact with mirror
Parallel swim	Fish swims upwards towards the mirror
Aggressive	
Chase	Fish quickly darts towards mirror
Bite	Fish opens jaw and bites mirror
Ram	Fish makes forceful contact with the mouth region to the mirror, often repetitively, but jaws remain closed
Head shake	Fish thrashes head from left to right repeatedly
Puffed throat	Fish flares out its operculum and lowers its jaw
Aggressive posture	
Lateral display	Fish presents its lateral aspect to the mirror while extending its unpaired fins
Pseudo-mouth fight	Back-and-forth movement occurs facing the mirror, as if fish will mouth fight, but no contact is made
Hook/J display	Fish swims towards the mirror, bites or rams it, then turns away and quivers
Submissive	
Submissive posture	Fish raises its head towards the mirror and lowers its tail
Submissive display	Fish in submissive posture but with a quivering tail
Flee	Fish quickly swims away from mirror

recordings (these periods of white noise went from milliseconds to minutes) and plotting their frequencies as a histogram. Any discontinuity would be indicative of how many pulses constitute a typical call. Lastly, the pulse rate can be defined as the function of the number of pulses per call duration.

The aforementioned temporal parameters were measured on the oscillogram in the same preset window size and settings, whereas peak frequency was quantified with the spectrogram (Hann, FFT size 256 samples, filter bandwidth 270 Hz, with a 50% overlap). All measurements were made in Raven Pro 1.5 sound analysis software, commonly employed in animal communication research (Bioacoustics Research Program, 2014). Analyses were performed in R 3.2.3 with model assumptions of normality and homogeneity of variance of the data initially confirmed (R Core Team, 2015).

Results

Of the seven (four males and three females) out of 10 *N. pulcher* that produced sound in the field experiment at Lake Tanganyika, there were a total of

40 pulses recorded (mean \pm SD; 5.7 ± 7.1 pulses/fish). These were short, broadband click sounds with pulse duration of 1.5 ± 0.5 ms and pulse peak frequency of 12008.0 ± 8312.8 Hz. In the laboratory setting, six (four males and two females) out of 10 *N. pulcher* emitted broadband click sounds of similar frequency (12938.7 ± 3494.0 Hz), but of longer duration (12.0 ± 3.4 ms; Wilcoxon-Mann-Whitney test, two-tail P value = 0.001). It should be noted that the fish, which did not produce sounds, do not necessarily reflect truly silent fish, rather the proportion of individuals that did not vocalise during the time allotted for recording. Importantly, the short, broadband high-frequency sounds described here were only detected when *N. pulcher* were present in the concrete pond or the glass aquarium, and never recorded on an empty setup or on sound recordings of the haplochromine cichlid *Astatoreochromis alluaudi* Pellegrin, 1904 done in the same experimental aquarium (Spinks, unpublished results).

A total of 92 pulses (14.8 ± 11.5 pulses/fish) produced by six individuals were measured in the laboratory setup (Table 2). Sound production was associated with all recorded behaviours, both aggressive and submissive, although five doubled-pulsed




Fig. 2 Oscillogram and spectrogram of a sound produced by *N. pulcher*. The oscillogram (A) presents the waveform of the pulses in time versus amplitude, while the spectrogram (B) shows how the frequency of the pulses changes over time, the colour indicates the relative amplitude. Here, the aforementioned temporal parameters; call duration (*a*), pulse duration (*b*) and interpulse interval (*c*) are illustratively defined. This double-pulsed call was made by a male in the laboratory experiments that concurrently exhibited an aggressive lateral display just after a series of rams and bites to the mirror

calls from two fish (one male and one female) were also recorded without concurrent visual display, when both fish were motionless (Table 2). This particular female had produced sound with behavioural displays, however, paused displaying for a couple of minutes and continued to call and then began displaying again. The male on the other hand did not display once, when he performed a few exploratory swims of the aquarium and then stayed in the corner of the aquarium calling out the rest of the recording. These sounds did not come from background or incidental noise and were similar to the other acoustic signals produced during displays (Table 2).

Inspection of interpulse duration frequency revealed that the majority of pulses were produced less than 0.4 s apart (Fig. 3). Pulses closer than 0.4 s were then considered part of one call, and on average, 2 pulses were produced per call (Additional Files 1–3; note similarity except for variation in temporal

properties). When this double-pulse call occurred, often the first pulse had a dominant frequency between 7000 Hz and 15000 Hz and the second pulse peaked higher, above 17,000 Hz (Fig. 2). Since the minimum resonance frequency of the aquarium ($\sim 4,000$ Hz) was much lower than the dominant frequency of *N. pulcher* sounds ($\sim 12,000$ Hz, Table 2), according to (Akamatsu et al., 2002) the resonance distortion in the aquarium should be minimal.

While the standard two-pulsed call was found in both sexes, male *N. pulcher* seemed to produce longer pulses than females (Table 3). Given the small number of vocal fish these differences indicate possible sex differences worth further examination.

Discussion

Sound production by Princess cichlids


Multimodal communication is expected in socially complex species (Freeberg et al., 2012) that live in unstable environments (Munoz & Blumstein, 2012). In this study, we report the production of short-duration, broadband click sounds often associated with a visual display by the cooperatively breeding Princess cichlid, *Neolamprologus pulcher*. Our analyses confirm recent findings that this species does not produce the low-frequency sounds common to many other cichlids and fish species in general (Pisanski et al., 2015), for which we suggest the term “low frequency silencing”. However, we found strong evidence for deliberate production of high-frequency double-pulse calls by *N. pulcher*. In our field and laboratory experiments, we found that both males and females produce broadband high-frequency sounds (above 5 kHz, average ~ 12 kHz) most often in an agonistic context induced by mirrors.

Broadband high-frequency sound production has long been reported in cichlids, including in species from Lake Tanganyika (e.g. Myrberg, Jr. et al., 1965; Nelissen, 1978), southern African *Oreochromis mossambicus* (Peters, 1852) (Lanzing, 1974) and more recently in West African *Pelmatolapia* (*Tilapia*) *mariae* (Boulenger, 1899) (Kottege et al., 2015). Reported peak frequencies of species from Lake Tanganyika are similarly high (above 5 kHz, often higher than 20 kHz), but temporal characteristics differ substantially among species. *Neolamprologus*

Table 2 Parameters (mean \pm SD) of the acoustic signals associated with and without a typical *Neolamprologus pulcher* social behaviour

	No. fish	Total pulses	Pulse duration (ms)	Pulse peak frequency (Hz)	Total calls	Call duration (ms)	Pulses per call
With behaviour	5	82	11.5 \pm 3.5	12280.5 \pm 3740.3	43	896.0 \pm 804.4	2.0 \pm 0.7
Without behaviour	2	10	13.2 \pm 2.8	13992.2 \pm 1889.3	5	294.4 \pm 324.0	2.0 \pm 0.0
Pooled	6	92	12.0 \pm 3.4	12938.7 \pm 3494.0	48	836.0 \pm 733.7	2.0 \pm 0.7

One fish emitted sound both with and without behaviour, therefore the pulses for each were calculated separately, except when pooled

**Fig. 3** Histogram of interpulse duration frequency. The majority of pulses within a call are shortly separated by less than 0.4 s

pulcher produces a distinct double-pulse clicking call, while others (*Astatotilapia burtoni* (Günther, 1894), *Simochromis diagramma* (Günther, 1893) and multiple *Tropheus* spp. Boulenger, 1898) produce a creaking or chewing multi-pulsed call (Nelissen, 1978). The recently reported short-duration broadband sound produced by *P. mariae* is very similar to the one described here for *N. pulcher*, except that it is single-pulsed (Kottege et al., 2015). The short pulses of sound and high frequency in *N. pulcher* point towards a stridulatory mechanism of sound production. It has

been suggested that African cichlids may produce sound by rubbing together the teeth on their pharyngeal jaws (Rice & Lobel, 2004), although this mechanism is yet to be confirmed. Fine & Parmentier (2015) suggest that stridulatory mechanisms should contain a wide range of frequencies, such as the broadband sound produced by *N. pulcher*.

The sounds recorded in this study were often produced in association with an aggressive visual display, but interestingly also in submissive displays. Importantly, since fish also produced sound with similar characteristics without an associated behaviour, we can infer that sound production is not a sole by-product of a visual display but instead can be generated independently. By examining both wild and captive fish, we could also exclude any effect of captivity and captive breeding on “low frequency silencing” in *N. pulcher*. The evolutionary reasons for loss of low frequency sounds are still unclear.

Acoustic differences between and within wild and captive individuals

Both wild and captive individuals generate characteristic high-frequency clicks, but pulses of *N. pulcher* in the laboratory recordings were longer in duration compared to the field recordings (one order of

Table 3 Sex differences in the parameters (mean \pm SD) of the acoustic signals of *Neolamprologus pulcher* in the laboratory experiments

	No. fish	Total pulses	Pulse duration (ms)	Pulse peak frequency (Hz)	Total calls	Call duration (ms)	Pulses per call
Male	4	64	14.1 \pm 2.1	12710.0 \pm 4303.6	36	918.0 \pm 770.4	1.8 \pm 0.1
Female	2	28	8.5 \pm 1.5	13396.2 \pm 2201.8	12	669.6 \pm 910.2	2.3 \pm 1.4
Pooled	6	92	12.0 \pm 3.4	12938.7 \pm 3494.0	48	836.0 \pm 733.7	2.0 \pm 0.7

All sounds produced were taken into account, both with and without a typical social behaviour

magnitude on average). Interestingly, male and female *N. pulcher* seem to differ in temporal parameters, although a more thorough investigation into the apparent temporal and sex differences would be required. Nevertheless cichlid acoustic studies have shown variation in pulse duration between closely related species, suggesting it is evolutionarily labile: mean pulse duration in *Oreochromis mossambicus* is 150 ms, compared to 10 ms in *Oreochromis niloticus* (Amorim et al., 2003; Longrie et al., 2008), and species in the genus *Maylandia* show 2- to 3-fold differences in mean pulse duration (Danley et al., 2012). Furthermore, context- and sex-specific differences have been reported in *Maylandia* (*'Pseudotropheus'*) *zebra* (Simões et al., 2008), and intra-individual variation in sound duration and pulse rate in response to motivation has been demonstrated in three distantly related cichlid species (Myrberg, Jr. et al., 1965). It is thus possible that noisier captive conditions have induced changes on labile temporal properties of *N. pulcher* sounds (pulse duration or period) in a similar way that environmental noise has impacted call duration and rate in Cope's grey treefrog, *Hyla chrysoscelis* Cope, 1880 (Love & Bee, 2010) or song amplitude in common blackbird, *Turdus merula* Linnaeus, 1,758 and other birds (Nemeth et al., 2013).

Significance of high-pitch sounds

Reports of low (i.e. below 2–3 kHz)-frequency sounds in cichlid fishes have been dominating the literature in recent years. This has likely both technical and biological explanations. On one hand, it is possible that sounds produced by cichlids in a reproductive context are mostly low frequency (e.g. Nelissen, 1978), while recording of narrower bandwidths or applying low-pass filters to raw data could account for masking of higher frequencies (e.g. Ripley & Lobel, 2004; Amorim et al., 2008; Longrie et al., 2008, 2009; Simões et al., 2008; Bertucci et al., 2012; Maruska et al., 2012; Pisanski et al., 2015). While unfiltered recordings of *N. pulcher* tested under different social contexts (Pisanski et al., 2015) did have a few intriguing high-frequency sounds with frequencies above 4 kHz, the temporal patterns were different from those reported here, and additional research is needed to determine if these were caused by incidental movement as opposed to originating from the fish as

purposeful communicatory sounds (S. Balshine, K. Pisanski and S. Marsh-Rollo personal communication). But perhaps the overarching reason relates to the expectation that fish are sensitive only to low-frequency sounds and cannot hear above a certain threshold (e.g. Heffner & Heffner, 1998), which would render such high-frequency sounds irrelevant for intraspecific communication. It is presently unclear whether *N. pulcher* can detect such high frequencies, as hearing sensitivities have not been studied in this species and those of the close-relative *N. brichardi* (Gante et al., 2016) have been investigated only in the range 100–2000 Hz (Ladich & Wysocki, 2003). Similarly, only low-frequency sensitivities (up to 1 or 2 kHz) were examined in other cichlids studied to date (Kenyon et al., 1998; Ripley et al., 2002; Smith et al., 2004). Nevertheless, evidence has been mounting that some fish species react to high-frequency sounds: for instance, behavioural studies indicate that cod *Gadus morhua* Linnaeus, 1758 can detect ultrasonic signals up to 38 kHz and the clupeid *Alosa sapidissima* (Wilson, 1811) of over 180 kHz, possibly allowing them to identify predatory echolocating cetaceans (Astrup & Møhl, 1993; Mann et al., 1997). Furthermore, new data indicate that some fish species might have multiple hearing maxima, as bimodal w-shaped sensitivity curves have been described in Malawian cichlids previously thought to have only a u-shaped sensitivity curve peaking at low frequencies (van Staaden et al., 2012).

Nelissen (1978) suggested that vocal complexity (measured as number of sound types) in six cichlid species from Lake Tanganyika varies inversely with number of colour patterns, such that different species would specialise along one of the two communication axes. Maruska et al. (2012) showed that acoustic signalling is an important sensory channel in multimodal courtship in the cichlid *A. burtoni*. Females responded to sounds even before seeing males (Maruska et al., 2012), which suggests that sounds could function as a long-distance attraction signal in the turbid waters of river deltas inhabited by this species. Sounds in the cooperative breeding *N. pulcher* could play a role in multimodal communication in an agonistic context and to maintain group cohesion. Since *N. pulcher* also produced sound in the confines of the shelter, it is possible that individuals can use acoustic signals when retreating to their shelter and other forms of communication are limited.

Importantly, high-frequency signals would also transmit more efficiently above the low-frequency background noise of the underwater world, particularly in windy conditions (van Staaden et al., 2012) or crowded fish neighbourhoods. These longer-range high-pitch sounds would allow communication among individuals belonging to different family groups, establishing a chorus across the colony.

While the ability of *N. pulcher* to hear in this high-frequency range is still to be determined, several hearing ‘specialists’ inhabiting Lake Tanganyika could be potential interspecific receivers of the acoustic signals generated by cichlids. Hearing ‘specialists’ that can detect sounds in the kHz generally have their swim bladder acoustically coupled to the inner ear and improved hearing in ‘generalists’ follows similar rules (Popper & Lu, 2000; Ladich & Schulz-Mirbach, 2016). ‘Specialists’ that can potentially hear higher frequency sounds and predate on cichlids include several catfish of the families Malapteruridae, Mochokidae, Claroteidae and Clariidae (Siluriformes). Other candidates would be the many species that lurk around *Neolamprologus* rocky habitat, such as spiny eels of the family Mastacembelidae (Synbranchiformes), tigerfish of the family Alestidae (Characiformes) and perches of the family Latidae (Perciformes).

Conclusion

We have shown that *N. pulcher* produces short-duration, broadband high-frequency (above 5 kHz, average ~12 kHz) double-pulsed calls. Sounds are most often produced jointly with aggressive or submissive visual displays, although both acoustic and visual signals can be produced in isolation. It is unclear whether the receiver of such sounds is intra- or inter-specific given our general lack of understanding of hearing sensitivities of fishes inhabiting Lake Tanganyika. In the event, that cichlids can hear such high-pitch sounds, an as of yet undescribed morphological adaptation transmitting vibrations of swim bladder walls to the inner ears is expected to exist. Non-visual sensory modalities in African cichlids may thus have a larger impact than originally expected and could be an important aspect in their adaptive radiation.

Acknowledgments We thank Lia Albergati and Benjamin Küng for assistance in the field and Miguel Vences for use of the recording equipment. We are grateful to Sigal Balshine, Katarzyna Pisanski and Susan E. Marsh-Rollo for sharing information on their raw data with us. This Project was funded by a travel Grant from the University of Basel to RKS, by Grants from the European Research Council (ERC; StG “CICHLID ~ X”) and the Swiss National Science Foundation (SNF) to WS, and by the “University of Basel Excellence Scholarships for Young Researchers” and “Novartis Excellence Scholarships for Life Sciences” to HFG.

References

- Akamatsu, T., T. Okumura, N. Novarini & H. Y. Yan, 2002. Empirical refinements applicable to the recording of fish sounds in small tanks. *The Journal of the Acoustical Society of America* 112: 3073–3082.
- Amorim, M. C. P., 2006. Diversity of sound production in fish. *Diversity* 1: 71–105.
- Amorim, M. C. P., P. J. Fonseca & V. C. Almada, 2003. Sound production during courtship and spawning of *Oreochromis mossambicus*: male–female and male–male interactions. *Journal of Fish Biology* 62: 658–672.
- Amorim, M. C. P., J. M. Simões, P. J. Fonseca & G. F. Turner, 2008. Species differences in courtship acoustic signals among five Lake Malawi cichlid species (*Pseudotropheus* spp.). *Journal of Fish Biology* 72: 1355–1368.
- Astrup, J. & B. Møhl, 1993. Detection of intense ultrasound by the cod *Gadus morhua*. *Journal of Experimental Biology* 182: 71–80.
- Aubin-Horth, N., J. K. Desjardins, Y. M. Martei, S. Balshine & H. A. Hofmann, 2007. Masculinized dominant females in a cooperatively breeding species. *Molecular Ecology* 16: 1349–1358.
- Bachmann, J. C., F. Cortesi, M. Hall, N. J. Marshall, W. Salzburger & H. F. Gante, 2016. Social selection maintains honesty of a dynamic visual signal in cichlid fish. *bioRxiv*. doi:10.1101/039313.
- Balshine-Earn, S. & A. Lotem, 1998. Individual recognition in a cooperatively breeding cichlid: evidence from video playback experiments. *Behaviour* 135: 369–386.
- Bertucci, F., J. Attia, M. Beauchaud & N. Mathevon, 2012. Sounds produced by the cichlid fish *Metriacrima zebra* allow reliable estimation of size and provide information on individual identity. *Journal of Fish Biology* 80: 752–766.
- Bradbury, J. W. & S. L. Vehrencamp, 2011. *Principles of Animal Communication*, 2nd ed. Sinauer Associates Inc, Sunderland, MA.
- Buchner, A. S., K. A. Sloman & S. Balshine, 2004. The physiological effects of social status in the cooperatively breeding cichlid *Neolamprologus pulcher*. *Journal of Fish Biology* 65: 1080–1095.
- Danley, P. D., M. Husemann & J. Chetta, 2012. Acoustic diversity in Lake Malawi’s rock-dwelling cichlids. *Environmental Biology of Fishes* 93: 23–30.
- Desjardins, J. K., K. A. Stiver, J. L. Fitzpatrick, N. Milligan, G. J. Van Der Kraak & S. Balshine, 2008. Sex and status in a

- cooperative breeding fish: behavior and androgens. *Behavioral Ecology and Sociobiology* 62: 785–794.
- Dijkstra, P. D., S. M. Schaafsma, H. A. Hofmann & T. G. G. Groothuis, 2012. “Winner effect” without winning: unresolved social conflicts increase the probability of winning a subsequent contest in a cichlid fish. *Physiology & Behavior* 105: 489–492.
- Fine, M. L. & E. Parmentier, 2015. Mechanisms of fish sound production. In Ladich, F. (ed.), *Sound Communication in Fishes*. Springer Vienna, Wien: 77–126.
- Freeberg, T. M., R. I. M. Dunbar & T. J. Ord, 2012. Social complexity as a proximate and ultimate factor in communicative complexity. *Philosophical Transactions of the Royal Society B: Biological Sciences* 367: 1785–1801.
- Frostman, P. & P. T. Sherman, 2004. Behavioral response to familiar and unfamiliar neighbors in a territorial cichlid, *Neolamprologus pulcher*. *Ichthyological Research* 51: 8–10.
- Gante, H. F. & W. Salzburger, 2012. Evolution: cichlid models on the runaway to speciation. *Current Biology* 22: R956–R958.
- Gante, H. F., M. Matschiner, M. Malmström, K. S. Jakobsen, S. Jentoft, & W. Salzburger, (2016). Genomics of speciation and introgression in Princess cichlid fishes from Lake Tanganyika. *Molecular Ecology*. doi:10.1111/mec.13767.
- Heffner, H. E. & R. S. Heffner, 1998. Hearing. In Greenberg, G. & M. M. Haraway (eds), *Comparative Psychology, A Handbook*. Garland, New York: 290–303.
- Kenyon, T. N., F. Ladich & H. Y. Yan, 1998. A comparative study of hearing ability in fishes: the auditory brainstem response approach. *Journal of Comparative Physiology A* 182: 307–318.
- Kohda, M., L. A. Jordan, T. Hotta, N. Kosaka, K. Karino, H. Tanaka, M. Taniyama & T. Takeyama, 2015. Facial recognition in a group-living cichlid fish. *PLoS One* 10: 1–12.
- Kottege, N., R. Jurdak, F. Kroon & D. Jones, 2015. Automated detection of broadband clicks of freshwater fish using spectro-temporal features. *The Journal of the Acoustical Society of America* 137: 2502–2511.
- Ladich, F., 1990. Vocalization during agonistic behaviour in *Cottus gobio* L. (Cottidae): an acoustic threat display. *Ethology* 84: 193–201.
- Ladich, F., 1997. Agonistic behaviour and significance of sounds in vocalizing fish. *Marine and Freshwater Behaviour and Physiology* 29: 87–108.
- Ladich, F. & R. R. Fay, 2013. Auditory evoked potential audiometry in fish. *Reviews in Fish Biology and Fisheries* 23: 317–364.
- Ladich, F. & T. Schulz-Mirbach, 2013. Hearing in cichlid fishes under noise conditions. *PLoS ONE* 8: e57588.
- Ladich, F. & T. Schulz-Mirbach, 2016. Diversity in fish auditory systems: one of the riddles of sensory biology. *Frontiers in Ecology and Evolution* 4: 1–26.
- Ladich, F. & L. E. Wysocki, 2003. How does tripus extirpation affect auditory sensitivity in goldfish? *Hearing Research* 182: 119–129.
- Lanzing, W. J. R., 1974. Sound production in the cichlid *Tilapia mossambica* Peters. *Journal of Fish Biology* 6: 341–347.
- Le Vin, A. L., B. K. Mable & K. E. Arnold, 2010. Kin recognition via phenotype matching in a cooperatively breeding cichlid, *Neolamprologus pulcher*. *Animal Behaviour* 79: 1109–1114.
- Lobel, P. S., I. M. Kaatz & A. N. Rice, 2010. Acoustical behavior of coral reef fishes. In Cole, K. (ed.), *Reproduction and Sexuality in Marine Fishes*. University of California Press, Berkeley, CA: 307–386.
- Longrie, N., M. L. Fine & E. Parmentier, 2008. Innate sound production in the cichlid *Oreochromis niloticus*. *Journal of Zoology* 275: 413–417.
- Longrie, N., S. Van Wassenbergh, P. Vandewalle, Q. Mauguit & E. Parmentier, 2009. Potential mechanism of sound production in *Oreochromis niloticus* (Cichlidae). *The Journal of Experimental Biology* 212: 3395–3402.
- Love, E. K. & M. A. Bee, 2010. An experimental test of noise-dependent voice amplitude regulation in Cope’s grey treefrog, *Hyla chrysoscelis*. *Animal Behaviour* 80: 509–515.
- Lugli, M., 2015. Habitat acoustics and the low-frequency communication of shallow water fishes. In Ladich, F. (ed.), *Sound Communication in Fishes*. Springer Vienna, Wien: 175–206.
- Lythgoe, J. N. & J. C. Partridge, 1991. The modelling of optimal visual pigments of dichromatic teleosts in green coastal waters. *Vision Research* 31: 361–371.
- Mann, D. A., Z. Lu & A. N. Popper, 1997. A clupeid fish can detect ultrasound. *Nature* 389: 341.
- Maruska, K. P. & R. D. Fernald, 2013. Social regulation of male reproductive plasticity in an African cichlid fish. *Integrative and Comparative Biology* 53: 938–950.
- Maruska, K. P., U. S. Ung & R. D. Fernald, 2012. The African cichlid fish *Astatotilapia burtoni* uses acoustic communication for reproduction: sound production, hearing, and behavioral significance. *PLoS One* 7: 1–13.
- Munoz, N. E. & D. T. Blumstein, 2012. Multisensory perception in uncertain environments. *Behavioral Ecology* 23: 457–462.
- Muschick, M., A. Indermaur & W. Salzburger, 2012. Convergent evolution within an adaptive radiation of cichlid fishes. *Current Biology* 22: 2362–2368.
- Myrberg Jr., A. A., E. Kramer & P. Heinecke, 1965. Sound production by cichlid fishes. *Science* 149: 555–558.
- Nelissen, M. H. J., 1978. Sound production by some Tanganyikan cichlid fishes and a hypothesis for the evolution of their communication mechanisms. *Behaviour* 64: 137–147.
- Nemeth, E., N. Pieretti, S. A. Zollinger, N. Geberzahn, J. Pardecke, A. C. Miranda & H. Brumm, 2013. Bird song and anthropogenic noise: vocal constraints may explain why birds sing higher-frequency songs in cities. *Proceedings of the Royal Society B: Biological Sciences* 280: 20122798.
- Parikh, V. N., T. Clement & R. D. Fernald, 2006. Physiological consequences of social descent: studies in *Astatotilapia burtoni*. *Journal of Endocrinology* 190: 183–190.
- Pisanski, K., S. E. Marsh-Rollo & S. Balshine, 2015. Courting and fighting quietly: a lack of acoustic signals in a cooperative Tanganyikan cichlid fish. *Hydrobiologia* 748: 87–97.
- Popper, A. N. & Z. Lu, 2000. Structure–function relationships in fish otolith organs. *Fisheries Research* 46: 15–25.
- R Core Team, 2015. R: A Language and Environment for Statistical Computing. Austria, Vienna.

- Radford, C. A., S. Ghazali, A. G. Jeffs & J. C. Montgomery, 2015. Vocalisations of the bigeye *Pempheris adspersa*: characteristics, source level and active space. *Journal of Experimental Biology* 218: 940–948.
- Rice, A. N. & P. S. Lobel, 2004. The pharyngeal jaw apparatus of the Cichlidae and Pomacentridae: function in feeding and sound production. *Reviews in Fish Biology and Fisheries* 13: 433–444.
- Ripley, J. L. & P. S. Lobel, 2004. Correlation of acoustic and visual signals in the cichlid fish, *Tramitichromis intermedius*. *Environmental Biology of Fishes* 71: 389–394.
- Ripley, J. L., P. S. Lobel & H. Y. Yan, 2002. Correlation of sound production with hearing sensitivity in the Lake Malawi cichlid *Tramitichromis intermedius*. *Bioacoustics* 12: 238–240.
- Rowland, W. J., 1999. Studying visual cues in fish behavior: a review of ethological techniques. *Environmental Biology of Fishes* 56: 285–305.
- Rumyantsev, S. L., S. Sawyer, N. Pala, M. S. Shur, Y. Bilenko, J. P. Zhang, X. Hu, A. Lunev, J. Deng & R. Gaska, 2005. Low frequency noise of light emitting diodes. *Proc SPIE Noise Dev Circ III* 5844: 75–85.
- Salzburger, W., 2009. The interaction of sexually and naturally selected traits in the adaptive radiations of cichlid fishes. *Molecular Ecology* 18: 169–185.
- Salzburger, W., B. Van Bocxlaer & A. S. Cohen, 2014. Ecology and evolution of the African Great Lakes and their faunas. *Annual Review of Ecology, Evolution, and Systematics* 45: 519–545.
- Santos, M. E. & W. Salzburger, 2012. Evolution. How cichlids diversify. *Science (New York, N.Y.)* 338: 619–621.
- Simões, J. M., I. G. Duarte, P. J. Fonseca, G. F. Turner & M. C. Amorim, 2008. Courtship and agonistic sounds by the cichlid fish *Pseudotropheus zebra*. *The Journal of the Acoustical Society of America* 124: 1332–1338.
- Smith, M. E., A. S. Kane & A. N. Popper, 2004. Acoustical stress and hearing sensitivity in fishes: does the linear threshold shift hypothesis hold water? *The Journal of Experimental Biology* 207: 3591–3602.
- Sopinka, N. M., J. L. Fitzpatrick, J. K. Desjardins, K. A. Stiver, S. E. Marsh-Rollo & S. Balshine, 2009. Liver size reveals social status in the African cichlid *Neolamprologus pulcher*. *Journal of Fish Biology* 75: 1–16.
- Stevens, M., 2013. *Sensory Ecology, Behaviour, and Evolution*. Oxford University Press, Oxford.
- van Oosterom, L., J. C. Montgomery, A. G. Jeffs & C. A. Radford, 2016. Evidence for contact calls in fish: conspecific vocalisations and ambient soundscape influence group cohesion in a nocturnal species. *Scientific Reports Nature Publishing Group* 6: 1–8.
- van Staaden, M., L. E. Wysocki & F. Ladich, 2012. Ecoacoustical constraints shape sound communication in Lake Malawi Cichlids [abstract]. *Bioacoustics* 21: 84.
- Verzijden, M. N., J. Van Heusden, N. Bouton, F. Witte, C. Ten Cate & H. Slabbekoorn, 2010. Sounds of male Lake Victoria cichlids vary within and between species and affect female mate preferences. *Behavioral Ecology* 21: 548–555.
- Wagner, C. E., L. J. Harmon & O. Seehausen, 2012. Ecological opportunity and sexual selection together predict adaptive radiation. *Nature Nature Publishing Group* 487: 366–369.
- White, S. A., T. Nguyen & R. D. Fernald, 2002. Social regulation of gonadotropin-releasing hormone. *The Journal of Experimental Biology* 205: 2567–2581.
- Wong, M. & S. Balshine, 2011. The evolution of cooperative breeding in the African cichlid fish, *Neolamprologus pulcher*. *Biological Reviews of the Cambridge Philosophical Society* 86: 511–530.
- Zöttl, M., D. Heg, N. Chervet & M. Taborsky, 2013. Kinship reduces alloparental care in cooperative cichlids where helpers pay-to-stay. *Nature Communications* 4: 1341.